

Cities and the Environment (CATE)

Volume 3 Issue 1 *MillionTreesNYC*, *Green Infrastructure*, and *Urban Ecology: A Research Symposium*

Article 25

December 2010

Using Phytotechnologies to Remediate Brownfields, Landfills, and Other Urban Areas

Ronald S. Zalesny Jr U.S. Forest Service, Northern Research Station, Institute for Applied Ecosystem Studies, rzalesny@fs.fed.us

Jill A. Zalesny

U.S. Forest Service, Northern Research Station, Institute for Applied Ecosystem Studies, jill.zalesny@wisconsin.gov

Follow this and additional works at: https://digitalcommons.lmu.edu/cate

Recommended Citation

Zalesny, Ronald S. Jr and Zalesny, Jill A. (2010) "Using Phytotechnologies to Remediate Brownfields, Landfills, and Other Urban Areas," *Cities and the Environment (CATE)*: Vol. 3: Iss. 1, Article 25. Available at: https://digitalcommons.lmu.edu/cate/vol3/iss1/25

This Conference Poster is brought to you for free and open access by the Center for Urban Resilience at Digital Commons @ Loyola Marymount University and Loyola Law School. It has been accepted for inclusion in Cities and the Environment (CATE) by an authorized administrator of Digital Commons at Loyola Marymount University and Loyola Law School. For more information, please contact digitalcommons@lmu.edu.

Using Phytotechnologies to Remediate Brownfields, Landfills, and Other Urban Areas

Urban areas requiring remedial work has prompted the use of phytotechnologies to improve water quality, soil health, and biodiversity, as well as to achieve sustainable social and economic goals. Phytotechnologies directly use plants to clean up contaminated groundwater, soil, and sediment. While woody and herbaceous crops are candidates for such remediation systems, trees within the genera Populus (poplars, cottonwoods, aspens) and Salix (willows) are ideal given their fast growth, extensive root systems, and elevated rates of photosynthesis and transpiration. The genetic diversity within these genera substantially increases the establishment and growth potential across heterogeneous sites. We have tested these trees for more than a decade across various sites and contaminants, which has resulted in developing phyto-recurrent selection, a method utilizing multiple testing cycles to evaluate, identify, and select favorable varieties with adequate genetic variation to guard against insect/disease outbreaks and changing edaphic conditions (especially those induced by contaminated soil and water) in the field. We will present information from our studies involving the selection and growth of trees performing well across variable site conditions (generalists) or sites with specific contaminant concerns (specialists) in order to enhance the success of phytotechnologies at brownfields, landfills, and other urban areas. This effort supports scientists and resource managers to acquire information that contributes to the deployment of systems that are ecologically and economically more sustainable versus traditional technologies, while the general public maintains environmental quality and protection of the natural resource base on which local and regional recreation, agriculture, and forestry depend.

Keywords

poplar, Populus, Salix, sustainability, waste management, wastewater irrigation, willow

Acknowledgements

n/a - included on the poster

Using Phytotechnologies to Remediate Brownfields, Landfills, & Other Urban Areas

Zalesny and Zalesny Rthy tote charalogie of Rendediation in the Urban Environment

U.S. Forest Service, Northern Research Station, Institute for Applied Ecosystem Studies, Rhinelander, WI 54501, USA




Copyright 2010 by the authors. All rights reserved. This work is licensed to the public under the Creative Commons Attribution License. Cities and the Environment is produced by the Urban Ecology Program, Department of Biology, Seaver College, Loyola Marymount University in cooperation with the USDA Forest Service. Published by The Berkeley Electronic Press (bepress). http://catejournal.org

Zalesny, R.S. Jr. and J.A. Zalesny. 2010. Using phytotechnologies to remediate brownfields, landfills, and other urban areas. Cities and the Environment. 3(1):poster 25. http://escholarship.bc.edu/cate/vol3/iss1/25.



Urban areas requiring remedial work has prompted the use of phytotechnologies to improve water quality, soil health, & biodiversity, as well as to achieve sustainable social & economic goals. Phytotechnologies directly use plants to clean up contaminated groundwater, soil, & sediment. An example of such technologies is phytoremediation, which is comprised of many processes.

Northern



While woody & herbaceous crops are candidates for such remediation systems, trees within the genera *Populus* (poplars, cottonwoods, aspens) & *Salix* (willows) are ideal given their elevated rates of photosynthesis and transpiration (A), fast growth (B), & extensive root systems (C). These trees are often known as "workhorses" towards remediation of contaminated areas.

The genetic diversity within these genera


Examples of phytoremediation projects conducted at the Institute for Applied Ecosystem Studies include:

substantially increases the establishment & growth potential across heterogeneous sites.

- · Landfill leachate / effluent
- · Municipal wastewater
- · Papermill fibercake effluent
- Polychlorinated biphenyls (PCB's)
- · Petroleum hydrocarbons
- Nitrates / fertilizer residues
- Ammonia
- Heavy metals













OBJECTIVES

We have tested these trees for more than a decade across various sites & contaminants, which has resulted in developing **phyto-recurrent selection**, a method utilizing multiple testing cycles to evaluate, identify, & select favorable varieties with adequate genetic variation to guard against insect/disease outbreaks & changing edaphic conditions (especially those induced by contaminated soil & water) in the field. The figure below is an example for phytoremediation of landfill leachate.



REMEDIATION & RESTORATION

When moving towards remediation of contaminated sites, workhorse species are needed to achieve positive ecosystem function (i.e., plant biomass) while maintaining species diversity & complexity. Once remediation is successful, restoration is more likely to be attainable. These basic principles of the continuum between complete ecosystem degradation & pre-disturbance condition (i.e., restoration) are depicted below, with brownfields & landfills shown as examples that follow.



Published by Digital Commons at Loyola Marymoght Thiversity and Loyola Law School, 2010


BROWNFIELDS

Abandoned or underused industrial & commercial facilities available for re-use, where expansion or redevelopment may be complicated by real or perceived environmental contaminations. Typically these sites are limited to reclamation, but remediation may be possible in some instances.



LANDFILL

Typically remediation & restoration is possible on these sites.



OTHER URBAN AREAS

Examples includes railyards, shipyards, & harbors.

PRACTICAL IMPLICATIONS

Cost-effective phytotechnologies are available to assist researchers, resource managers, regulators, etc. to improve environmental conditions while protecting human health.

Phyto-recurrent selection can help researchers & resource managers move closer to obtaining long-term (i.e., rotation age) remediation data from brownfields, landfills, & other urban areas.

Well-developed experimental designs & associated precision levels support estimation of quantitative genetic data that lead to recommendations of generalist genotypes that perform well over broad contaminant ranges or specialist genotypes that perform well for specific contaminants.

This effort supports scientists & resource managers to acquire information that contributes to the deployment of phytotechnologies that are ecologically & economically more sustainable versus traditional systems, while the general public maintains environmental quality & protection of the natural resource base on which local & regional recreation, agriculture, & forestry depend.

ACKNOWLEDGMENTS

R.S. Zalesny Jr. presented this information at the Million TreesNYC, Green Infrastructure and Urban Ecology: A Research Symposium, March 5-6, 2010, New York, NY. This research is funded by the U.S. Forest Service Northern Research Station – Research Work Unit RWU-NRS-13, and has been supported by Iowa State University

Department of Natural Resource Ecology and Management. Contributions of Edmund 1 Bauer, David Coyle, Richard Hall, Neil Nelson, and Adam Wiese are greatly appreciated.

