



Digital Commons@

Loyola Marymount University
LMU Loyola Law School

Mathematics, Statistics and Data Science
Faculty Works

Mathematics, Statistics and Data Science

9-2012

Forecasting the Effect of the Amethyst Initiative on College Drinking

Ben G. Fitzpatrick

Loyola Marymount University, ben.fitzpatrick@lmu.edu

Richard Scribner

Louisiana State University

Azmy S. Ackleh

University of Louisiana at Lafayette

Jawaid Rasul

Biomedware Incorporated

Geoffrey Jacquez

The University Of Michigan

See next page for additional authors

Follow this and additional works at: https://digitalcommons.lmu.edu/math_fac



Part of the [Mathematics Commons](#)

Digital Commons @ LMU & LLS Citation

Fitzpatrick, Ben G.; Scribner, Richard; Ackleh, Azmy S.; Rasul, Jawaid; Jacquez, Geoffrey; Simonsen, Neal; and Rommel, Robert, "Forecasting the Effect of the Amethyst Initiative on College Drinking" (2012).

Mathematics, Statistics and Data Science Faculty Works. 96.

https://digitalcommons.lmu.edu/math_fac/96

This Article - post-print is brought to you for free and open access by the Mathematics, Statistics and Data Science at Digital Commons @ Loyola Marymount University and Loyola Law School. It has been accepted for inclusion in Mathematics, Statistics and Data Science Faculty Works by an authorized administrator of Digital Commons@Loyola Marymount University and Loyola Law School. For more information, please contact digitalcommons@lmu.edu.

Authors

Ben G. Fitzpatrick, Richard Scribner, Azmy S. Ackleh, Jawaid Rasul, Geoffrey Jacquez, Neal Simonsen, and Robert Rommel



Published in final edited form as:

Alcohol Clin Exp Res. 2012 September ; 36(9): 1608–1613. doi:10.1111/j.1530-0277.2012.01765.x.

Forecasting the Effect of the Amethyst Initiative on College Drinking

Ben G. Fitzpatrick, PhD^{1,2}, Richard Scribner, MD, MPH³, Azmy S. Ackleh, PhD⁴, Jawaid Rasul, PhD⁵, Geoffrey Jacquez, PhD^{5,6}, Neal Simonsen, PhD³, and Robert Rommel, PhD⁵

¹Loyola Marymount University, 1 LMU Drive, Los Angeles, CA 90045

²Tempest Technologies, 8939 S. Sepulveda Blvd, Suite 506, Los Angeles, CA 90045

³Louisiana State University School of Public Health, 1615 Poydras St., Room 1535, New Orleans, LA 70112

⁴University of Louisiana at Lafayette, P.O. Box 41010, Lafayette, LA 70504-1010

⁵Biomedware Incorporated, 3526 W. Liberty, Suite 100, Ann Arbor, MI 48103

⁶The University of Michigan School of Public Health, 1415 Washington Heights, Ann Arbor, MI 48109-2029

Abstract

Background—A number of college presidents have endorsed the Amethyst Initiative, a call to consider lowering the minimum legal drinking age (MLDA). Our objective is to forecast the effect of the Amethyst Initiative on college drinking.

Methods—A system model of college drinking simulates MLDA changes through (1) a decrease in heavy episodic drinking (HED) due to the lower likelihood of students drinking in unsupervised settings where they model irresponsible drinking (misperception), and (2) an increase in overall drinking among currently underage students due to increased social availability of alcohol (wetness).

Results—For the proportion of HEDs on campus, effects of large decreases in misperception of responsible drinking behavior were more than offset by modest increases in wetness.

Conclusions—For the effect of lowering the MLDA, it appears that increases in social availability of alcohol have a stronger impact on drinking behavior than decreases in misperceptions.

Keywords

Amethyst Initiative; Modeling; College; Heavy Episodic Drinking; Misperception

Introduction

College drinking is one of the most significant and complex public health problems today. Heavy drinking among college students remains a pervasive problem that places students at considerable risk for a variety of negative outcomes, including date rape, academic problems, traffic accidents, and health problems (Hingson, Heeren, Winter, and Wechsler, 2005; Wechsler and Nelson, 2008). Alcohol use is embedded in the college lifestyle, resulting in enormous social, economic, and health consequences among some of the nation's finest students (Task Force on College Drinking, 2002). Heavy episodic drinking is generally conducted in private, among peers, and college students engage in the behavior in much higher proportions than do other young adults (Schulenberg, et al., 2001; Carey, Scott-Sheldon, Carey, and DeMartini, 2007; Substance Abuse and Mental Health Services Administration, 2006; Timberlake, et al., 2007).

Interventions to reduce the negative outcomes associated with college drinking have been mixed. For example, reeducation programs targeting misperception of drinking norms remain a popular intervention. Social norms researchers have found that college students routinely misperceive the level of alcohol use among their peers (Baer, Stacy, and Larimer, 1991; Perkins, Haines, and Rice, 1991) and liberal perceptions of social norms for peer drinking are consistently shown to be strong predictors of alcohol use among college students (Baer, Stacy, and Larimer, 1991; Perkins, Haines, and Rice, 1991; Reis and Riley, 2000; Baer and Carney, 1993, Perkins, et al., 1999; Babor, Aguirre-Molina, Marlatt, and Clayton, 1999; Thombs, Wolcott, and Farkash, 1997). Rather than address the population level factors that lead to misperception of social norms in the first place, Social Norms Marketing (SNM) interventions attempt to reeducate students by correcting their misperceptions of their peers' behavior in hopes of changing individual behavior. The results of these interventions have been equivocal (Thombs, Wolcott, and Farkash, 1997; Haines and Spear, 1996; Werch, et al., 2000; Wechsler, et al., 2003; Toomey, Lenk, and Wagenaar, 2007; DeJong, et al., 2006).

The limited effectiveness of college drinking interventions has led to calls to reexamine the MLDA. One of these, called the Amethyst Initiative, asks chancellors and presidents of universities and colleges across the country to sign on to a call asking elected officials to revisit the 21 year old drinking age. So far over 100 chancellors and presidents have signed on. The Amethyst Initiative statement argues that the 21 year old drinking age is not working. Underage students are legally prohibited from purchasing and possessing alcohol and the majority continue to drink. In addition, the statement argues the rampant flaunting of the drinking laws by students has led to a "culture of dangerous binge drinking" on many campuses. John McCardell, the original author of the initiative, has described the mechanism by which this culture has evolved. He argues that because of the 21 MLDA underage students are precluded from drinking in supervised settings (e.g., bars, school sponsored parties). As a result, they are more likely to drink in unsupervised settings (e.g., off campus parties) where there are fewer constraints on excessive drinking. He concludes that in these settings underage students who are new to drinking develop misperceptions of the normal drinking behavior and binge drinking tends to be viewed as normal (McCardell, 2008). This argument is consistent with literature that demonstrates misperception of

drinking norms predicts individual drinking (Baer, Stacy, and Larimar, 1991; Perkins, Haines, and Rice, 2005). Social norms theory argues that the effect of misperceptions is rooted in a psychological attribution process in which the individual tends to perceive the drinking actions of others as reflective of their individual drinking temperament and align their behavior accordingly (Perkins, 1997; Prentice and Miller, 1993), proponents suggest that the MLDA of 21 years of age is the problem. Critics of the Amethyst Initiative argue that lowering the MLDA will increase the availability of alcohol for both social and home consumption, and therefore increase drinking among the entire population with disastrous results (Babor, 2008).

Lowering the current MLDA represents an enormous social experiment with potentially major consequences. While there is considerable evidence indicating the harms associated with lowering the MLDA with regard to the general population (Wagenaar and Toomey, 2002), there is little in the way of observational evidence (Kypri, et al., 2006) to either support or oppose the specific hypotheses regarding student drinking behaviors embedded in the Amethyst initiative. A systems approach is one method of providing a forecast for the effects of a policy change prior to carrying out interventions based on that policy. Public health researchers are beginning to see the opportunities of moving from a purely inferential approach of experimental design and data analysis to a more mechanistic, systems approach (Homer and Hirsch, 2006). In particular, the social and economic cost of suboptimal policy decisions can potentially be mitigated by an increased understanding of the potential consequences that a systems model can provide.

Materials and Methods

We have developed a systems model, referred to as SimHED (Ackleh, et al., 2009; Scribner, et al., 2009), to simulate a college campus student population structured by drinking behavior and drinking age. The model, a continuous dynamical systems compartmental model derived using epidemiological reasoning, is provided in detail in the Appendix. The compartmental structure involves two levels of structure in age, namely underage (U) and legal age (L), as well as four drinking styles associated with college drinking, namely abstainers (1) social drinkers (2), problem drinkers (3), and heavy episodic drinkers or HEDs (4). Abstainers are defined as individuals who do not drink. Social drinkers are individuals who drink more frequently than abstainers but do not belong to the other two compartments. Problem drinkers are those who report at least two out of four indicators of problem drinking based on the CAGE instrument (Mayfield, McLeod, and Hall, 1974), regardless of the amount or frequency of their drinking. Heavy episodic drinkers are individuals who consume more than five drinks in a single sitting on at least one occasion in the past two weeks. In this manner, the state of the system at any given time is defined by 8 numbers, $U_1, U_2, U_3, U_4, L_1, L_2, L_3, L_4$, which are the number of individuals in each of the four drinking style compartments for the underage drinkers (U_1, U_2, U_3, U_4) followed by the corresponding numbers for the legal age drinkers (L_1, L_2, L_3, L_4).

Individuals move from underage to legal age purely by aging. We assume that the aging process itself does not directly change the drinking style of the individual, so that transition from the underage to legal age compartments preserves the drinking style. The drinking

style transition model includes three types of parameters that control transfers between the four drinking style compartments: individual risk (Wechsler, Dowdall, Davenport, and Castillo, 1995; Presley, Meilman, and Leichter, 2002), social interactions (Reifman, Watson, and McCourt, 2002; McCabe, et al., 2005), and social norm misperception (Perkins, et al., 1999; Borsari and Carey, 2001).

The individual risk model handles transitions that depend only on individual factors (e.g., mood, developmental transitions, individual traits). This component of the system involves a fraction of individuals with a particular drinking style transitioning to a different drinking style over a period of time. These transitions are modeled by terms of the form $r_{ij}N_i$ (in which N represents U or L , r models the fraction of those individuals transitioning, and the subscript ij represents the transition out of drinking style i into drinking style j). Since the movement is from i to j , this term is a positive term in the j equation and a negative term in the i equation.

Social interaction transitions depend on individuals from two separate groups coming into contact with one another, much like an epidemiological model of disease transmission. That is, an individual with a certain type of drinking style convinces another individual with a different drinking style to change behavior. We model these transitions by terms of the form $s_{ij}N_iN_j$, proportional to the number of pairings available among the two different drinking styles. In this case, the movement may be in either direction, i.e., from i to j or vice versa. So, $s_{ij}N_iN_j$ represents a net movement between the two compartments i and j , and thus s_{ij} may be negative.

Social norms/misperception transitions occur due to perception of the level of a particular drinking behavior. These movements occur in two situations: when abstainers become light drinkers because they perceive an exaggeratedly large number of drinkers on campus, and when social drinkers become HEDs because they perceive an exaggeratedly large number of

HEDs. Situation a) is modeled by the term $n_{12}N_1M\left(\frac{\sum_{k=2}^4 U_k + L_k}{\sum_{k=1}^4 U_k + L_k}\right)$ and situation b) is

modeled by the terms $n_{i5}N_iM\left(\frac{U_4 + L_4}{\sum_{k=1}^4 U_k + L_k}\right)$, where i can be either 2 or 3. Again, N represents either U or L here, an underage (U) or legal age (L) drinking style, as the changes in drinking style are assumed to occur within an age group. The function M is the misperception function, modeling how badly the students overestimate the fraction of individuals undertaking the role model behavior (either the fraction of drinkers or the fraction of HEDs). Were M the identity function, student perception would be entirely accurate. Research in social norms suggests (Reis and Riley, 2000) that misperception is greatest when the model behavior is least prevalent and that misperception decreases as the model behavior increases. For example, in an analysis of the National College Health Assessment, 59.9% of students overestimate the drinking norms of their peers at parties by three or more drinks on campuses where abstinence is the norm. Where six drinks is the actual norm, 31.5% of students overestimate by three or more drinks (Perkins, Haines, and Rice, 2005). For these reasons we have chosen the functional form $M(x) = \sqrt{\varepsilon + (1 - \varepsilon)x^2}$ whose graph is illustrated in Figure 1.

The hyperparameter ε controls the level of misperception: as $\varepsilon \rightarrow 0$ the amount of overestimation goes to 0, and M becomes the identity function. We use the term “hyperparameter” here to distinguish from the basic transition rate parameters in the model and to emphasize that those rate parameters depend on ε .

The campus alcohol environment (i.e., level of campus wetness) is an additional hyperparameter, w , which modifies the transfer rates between compartments as a function of campus wetness. Each rate parameter r_{ij} , s_{ij} , n_{ij} depends linearly on the wetness. For example, $r_{23} = r_{23}^0(1 - w) + r_{23}^1w$, so that a completely “dry” campus, $w=0$, has rate parameter r_{23}^0 , and a completely “wet” campus, $w=1$, has rate parameter r_{23}^1 .

A version of the model without age structure has been successfully calibrated to survey data obtained from 32 campuses across the United States in the Social Norms Marketing Research Project, SNMRP (DeJong, et al., 2006; Ackleh, et al., 2009; Scribner, et al., 2009). We have reasonable estimates of the “wet” and “dry” rate parameter values, the wetness levels for each of the 32 campuses, and the misperception levels. Table 1 contains parameter estimates for the wet and dry rate parameters.

The wetness hyperparameters range from 0.05 to 0.55 in the 32 schools from the SNMRP data. Wetness correlates ($R^2 = 0.30$) with alcohol outlet density, a common measure of availability, but clearly other phenomena are involved, such as enforcement and campus social environment. However, lowering the MLDA would dramatically affect the effective alcohol outlet density for underage students. We estimated the misperception hyperparameters for the SNMRP campuses, and these values range from nearly 0 to 0.25.

We are interested in the exploration of the interplay between misperception and availability. In particular, the Amethyst Initiative's hypothesis is that a reduction in misperception and attendant improper role model choices will lead to a reduction in HED behavior. Toward that end, we consider a hypothetical college population behavior averaged over a ten year period, and we conduct a series of experiments. We assume that the wet and dry rate parameter values are the same for the under age and legal age groups; however, we take the legal age group to have higher wetness (greater availability) and lower misperception levels. We conducted a series of computer simulation experiments that we report here.

Results

We begin our exploration by considering a campus with medium wetness of 0.30, applicable to the legal age students, who also have a small amount of misperception ($\varepsilon=0.05$). In order to examine the effect of misperception, we consider a number of simulated “treatment” scenarios in which we assume a range of effects on the wetness and misperception parameters for the underage population and observe the resulting drinking behavior.

Our first simulation involves the assumption that the wetness hyperparameter for the underage population is the same as for the legal age ($w=0.30$), but that the misperception is at a higher level ($\varepsilon=0.25$). Our treatment is assumed to reduce the misperception from 0.25 down to 0.05, the level of the legal age students, with 100% treatment implementation

(Figure 2 upper panel). The x-axis in the upper panel Figure 2 denotes this linear change of misperception, where 0% implementation corresponds to $\varepsilon=0.25$ and 100% to $\varepsilon=0.05$. We can see that the reduction in Heavy Episodic Drinkers is relatively small. We should note that it is not clear that the legal age population would actually have less misperception than the underage population; however, such an assumption leads to a slightly conservative estimate on heavy episodic drinking.

A different approach might be to change the wetness parameter for the underage students. Our second simulation, in the lower panel of Figure 2, shows the fraction of HEDs in the legal and underage compartments as we reduce wetness in the underage group from 0.30 (0 percent treatment on the x-axis) down to 0.00 (100 percent treatment). Again, the legal age students have a small misperception parameter ($\varepsilon=0.05$) while the misperception of the underage students is at a high level ($\varepsilon=0.25$). One can see in the lower panel of Figure 2 that this treatment has a much stronger impact on the HED fraction than does the reduction of misperception.

Neither of these simulations captures the actual effect of the Amethyst Initiative's MLDA reduction. One might expect that wetness for the underage population is somewhat less than it is for the legal age population (exactly how much is of course a difficult matter to resolve) while the amount of misperception is greater for the underage population. We have conducted a number of simulations in which wetness and misperception for the underage population are changed simultaneously. As a first illustrative example, we continue with the hypothetical campus having wetness of 0.30 and misperception of 0.05 for the legal age students. We have also simulated a wet campus ($w=0.55$). We assume in this example that the underage population has a wetness parameter of half that of the legal age population, and a misperception parameter of 0.25, while the misperception parameter is at the high end of those we have inferred from SNMRP data. The treatment is to increase the underage wetness to that of the legal age group, while reducing the misperception to 0.05, so that at the end of the treatment, the legal age and underage students have the same parameters.

In Figure 3, we show the effect of simultaneously increasing the wetness and decreasing the misperception in the underage population. In each of the simulated experiments (both moderate and wet), the underage population goes from half the wetness of the legal population to fully as wet, while simultaneously going from high misperception (0.25) to the low level of the legal age population (0.05). In each of the three panels, the qualitative trend is the same: HED drinking among underage students is increased.

With any computer simulation model of a real-world phenomenon, but most especially with the highly challenging modeling problems of social systems, the prediction of events that are out of the scope of observation must be viewed with some skepticism. We have endeavored to calibrate and validate our model as accurately as possible (Ackleh, et al., 2009), and we have also conducted a large number of simulation studies to examine the dependence on assumptions about the legal age population, dropout and recruitment rates, and other parameters. Such sensitivity analyses of the model to its parameters are a necessary step for developing confidence in the resulting predictions. We have found across a wide spectrum

of such analyses that the structure of the basic findings presented here is remarkably consistent.

Discussion

Limitations

It is of course a risky exercise to attempt to predict possible behavior based on inferred parameterizations. Rather than viewing these results as quantitative predictions of the actual levels of drinking that will occur under an MLDA change, we prefer to interpret the model output in terms of the trends and joint behavior as wetness index and misperception are changed simultaneously. The inescapable conclusion is that the misperception must be very great among the underage population and very significantly reduced by allowing the underage population to drink in order to compensate for the increased availability of drinking venues to the underage population. Indeed, further studies are required to quantify with accuracy how these parameters might actually change in the presence of an MLDA reduction. We do, however, interpret our simulations to date as pessimistic for the Amethyst Initiative's proposal that an MLDA reduction will have beneficial consequences for college drinking.

Conclusions

The preliminary insights provided by this model suggest that a reduction in the MLDA may not produce the desired reduction in heavy episodic drinking that is the goal of the Amethyst Initiative's strategy, based on the Initiative's reasoning about why "21 is not working." (20). The analysis we have conducted suggests that effects of a reduction in misperception from the largest observed values to the lowest is overcome by a 25 percent increase in campus wetness. One might expect a much larger increase in wetness from the increased physical availability of alcohol associated with making the entire college population, rather than approximately half, to be of legal drinking age.

Acknowledgments

Support: This research was supported by National Institute on Alcohol Abuse and Alcoholism grant RO1 AA015573.

We dedicate this paper to the memory of our friend and colleague, Dr. Jawaid Rasul, who passed away shortly after the completion of this work.

Appendix

The eight-compartment set of equations is given by

$$\frac{dU_1}{dt} = -d_1U_1 - n_{12}f \left(\frac{\sum_{i=2}^4 (U_i + L_i)}{\sum_{i=1}^4 (U_i + L_i)} \right) U_1 - s_{12}U_1(U_2 + L_2) + r_{21}U_2 + r_{31}U_3 - \lambda_1U_1$$

$$\frac{dL_1}{dt} = -d_1 L_1 - n_{12} f \left(\frac{\sum_{i=2}^4 (U_i + L_i)}{\sum_{i=1}^4 (U_i + L_i)} \right) L_1 - s_{12} L_1 (U_2 + L_2) + r_{21} L_2 + r_{31} L_3 + \lambda_1 U_1$$

$$\begin{aligned} \frac{dU_2}{dt} = & -d_2 U_2 + n_{12} f \left(\frac{\sum_{i=2}^4 (U_i + L_i)}{\sum_{i=1}^4 (U_i + L_i)} \right) U_1 \\ & + s_{12} (U_2 + L_2) U_1 \\ & + s_{42} (U_2 + L_2) U_4 \\ & - n_{24} f \left(\frac{(U_4 + L_4)}{\sum_{i=1}^4 (U_i + L_i)} \right) U_2 \\ & - r_{21} U_2 - r_{23} U_2 - r_{24} U_2 + r_{42} U_4 - \lambda_2 U_2 \end{aligned}$$

$$\begin{aligned} \frac{dL_2}{dt} = & -d_2 L_2 + n_{12} f \left(\frac{\sum_{i=2}^4 (U_i + L_i)}{\sum_{i=1}^4 (U_i + L_i)} \right) L_1 \\ & + s_{12} (U_2 + L_2) L_1 \\ & + s_{42} (U_2 + L_2) L_4 \\ & - n_{24} f \left(\frac{(U_4 + L_4)}{\sum_{i=1}^4 (U_i + L_i)} \right) L_2 \\ & - r_{21} L_2 - r_{23} L_2 - r_{24} L_2 + r_{42} L_4 + \lambda_2 U_2 \end{aligned}$$

$$\frac{dU_3}{dt} = -d_3 U_3 - r_{31} U_3 + r_{23} U_2 + r_{43} U_4 - \lambda_3 U_3$$

$$\frac{dL_3}{dt} = -d_3 L_3 - r_{31} L_3 + r_{23} L_2 + r_{43} L_4 + \lambda_3 U_3$$

$$\frac{dU_4}{dt} = -d_4 U_4 - s_{42} U_4 (U_2 + L_2) + n_{24} f \left(\frac{(U_4 + L_4)}{\sum_{i=1}^4 (U_i + L_i)} \right) U_2 - r_{42} U_4 - r_{43} U_4 + r_{24} U_2 - \lambda_4 U_4$$

$$\frac{dL_4}{dt} = -d_4L_4 - s_{42}L_4(U_2 + L_2) + n_{24}f \left(\frac{(U_4 + L_4)}{\sum_{i=1}^4 (U_i + L_i)} \right) L_2 - r_{42}L_4 - r_{43}L_4 + r_{24}L_2 + \lambda_4U_4$$

References

- Ackleh AS, Fitzpatrick BG, Scribner R, Simonsen N, Thibodeaux J. Ecosystem Models of College Drinking: Parameter Estimation and Comparing Models to Data. *Mathematical and Computer Modelling*. 2009; 50:481–97. [PubMed: 20161275]
- Babor, T. Amethyst initiative versus animal house: What college presidents can learn from research on the Mediterranean drinking style. American Public Health Association Annual Meeting; San Diego, CA, USA. 2008.
- Babor TF, Aguirre-Molina M, Marlatt GA, Clayton R. Managing alcohol problems and risky drinking. *American Journal of Health Promotion*. 1999; 14(2):98–103. [PubMed: 10724728]
- Baer JS, Carney MM. Biases in the perceptions of the consequences of alcohol use among college students. *Journal of Studies on Alcohol and Drugs*. 1993; 54(1):54–60.
- Baer JS, Stacy A, Larimer M. Biases in the perception of drinking norms among college students. *Journal of Studies on Alcohol and Drugs*. 1991; 52(6):580–6.
- Borsari B, Carey KB. Peer influences on college drinking: a review of the research. *Journal of Substance Abuse*. 2001; 13(4):391–424. [PubMed: 11775073]
- Carey KB, Scott-Sheldon LA, Carey MP, DeMartini KS. Individual-level interventions to reduce college student drinking: a meta-analytic review. *Addictive Behaviors*. 2007; 32(11):2469–94. [PubMed: 17590277]
- DeJong W, Schneider SK, Towvim LG, Murphy MJ, Doerr EE, Simonsen NR, Mason K, Scribner RA. A multisite randomized trial of social norms marketing campaigns to reduce college student drinking. *Journal of Studies on Alcohol and Drugs*. 2006; 67(6):868–79.
- Haines M, Spear SF. Changing the perception of the norm: a strategy to decrease binge drinking among college students. *Journal of American College Health*. 1996; 45(3):134–40. [PubMed: 8952206]
- Hingson R, Heeren T, Winte M, Wechsler H. Magnitude of alcohol-related mortality and morbidity among U.S. college students ages 18-24: changes from 1998 to 2001. *Annual Review of Public Health*. 2005; 26:259–79.
- Homer JB, Hirsch GB. System dynamics modeling for public health: background and opportunities. *American Journal of Public Health*. 2006; 96(3):452–8. [PubMed: 16449591]
- Kypri K, Voas RB, Langley JD, Stephenson SCR, Begg DJ, Tippetts AS, Davie GS. Minimum purchasing age for alcohol and traffic crash injuries among 15- to 19- year-olds in New Zealand. *American Journal of Public Health*. 2006; 96(1):126–131. [PubMed: 16317197]
- Mayfield D, McLeod G, Hall P. The CAGE questionnaire: validation of a new alcoholism screening instrument. *American Journal of Psychiatry*. 1974; 131(10):1121–3. [PubMed: 4416585]
- McCabe SE, Schulenberg JE, Johnston LD, O'Malley PM, Bachman JG, Kloska DD. Selection and socialization effects of fraternities and sororities on US college student substance use: a multi-cohort national longitudinal study. *Addiction*. 2005; 100(4):512–24. [PubMed: 15784066]
- McCardell J. The drinking age. The status quo has bombed. *US News and World Report*. 2008; 145(6):11. [PubMed: 18831099]
- Perkins, HW. College student misperceptions of alcohol and other drug norms among peers; explaining causes, consequences and implications for prevention programs In: *The Higher Education Center for Alcohol and Other Drug Prevention, Designing Alcohol and Other Drug Prevention Programs in Higher Education*. Newton: US Department of Education; 1997.

- Perkins HW, Haines MP, Rice R. Misperceiving the college drinking norm and related problems: a nationwide study of exposure to prevention information, perceived norms and student alcohol misuse. *Journal of Studies on Alcohol and Drugs*. 2005; 66(4):470–8.
- Perkins HW, Meilman PW, Leichter JS, Cashin JR, Presley CA. Misperceptions of the norms for the frequency of alcohol and other drug use on college campuses. *Journal of American College Health*. 1999; 47(6):253–8. [PubMed: 10368559]
- Prentice DA, Miller DT. Pluralistic ignorance and alcohol use on campus: some consequences of misperceiving the social norm. *Journal of Personality and Social Psychology*. 1993; 64(2):243–56. [PubMed: 8433272]
- Presley CA, Meilman PW, Leichter JS. College factors that influence drinking. *Journal of Studies on Alcohol Supplement*. 2002; 14:82–90. [PubMed: 12022732]
- Reifman A, Watson WK, McCourt A. Social networks and college drinking: probing processes of social influence and selection. *Personality and Social Psychology Bulletin*. 2006; 32(6):820–32. [PubMed: 16648206]
- Reis J, Riley WL. Predictors of college students' alcohol consumption: implications for student education. *The Journal of Genetic Psychology*. 2000; 161(3):282–91. [PubMed: 10971907]
- Schulenberg J, Maggs JL, Long SW, Sher KJ, Gotham HJ, Baer JS, Kivlahan DR, Marlatt GA, Zucker RA. The problem of college drinking: insights from a developmental perspective. *Alcoholism: Clinical and Experimental Research*. 2001; 25(3):473–7.
- Scribner R, Ackleh A, Fitzpatrick B, Rommel R, Jacquez G, Simonsen N. A Systems Approach to College Drinking: Development of a Deterministic Model for Testing Alcohol Control Policies. *Journal of Studies on Alcohol and Drugs*. 2009; 70(5):805–21. [PubMed: 19737506]
- Substance Abuse and Mental Health Services Administration. Results from the 2005 national Survey on Drug Use and Health: National Findings. Rockville: DHHS; 2006. Office of Applied Studies, NSDUH Series H-30, DHHS Publication No. SMA 06-4194
- Task Force on College Drinking. Call to Action: Changing the Culture of College Drinking. Bethesda: National Institute of Alcohol Abuse and Alcoholism; 2002.
- Timberlake DS, Hopfer CJ, Rhee SH, Friedman NP, Haberstick BC, Lessem JM, Hewitt JK. College attendance and its effect on drinking behaviors in a longitudinal study of adolescents. *Alcoholism: Clinical and Experimental Research*. 2007; 31(6):1020–30.
- Thombs DL, Wolcott BJ, Farkash LG. Social context, perceived norms and drinking behavior in young people. *Journal of Substance Abuse*. 1997; 9:257–67. [PubMed: 9494953]
- Toomey TL, Lenk KM, Wagenaar AC. Environmental policies to reduce college drinking: an update of research findings. *Journal of Studies on Alcohol and Drugs*. 2007; 68(2):208–19. [PubMed: 17286339]
- Wagenaar AC, Toomey TL. Effects of Minimum Drinking Age Laws: Review and Analyses of the Literature from 1960 to 2000. *Journal of Studies on Alcohol and Drugs Supplement*. 2002; 14:206–225.
- Wechsler H, Dowdall GW, Davenport A, Castillo S. Correlates of college student binge drinking. *American Journal of Public Health*. 1995; 85(7):921–6. [PubMed: 7604914]
- Wechsler H, Nelson TF. What we have learned from the Harvard School Of Public Health College Alcohol Study: focusing attention on college student alcohol consumption and the environmental conditions that promote it. *Journal of Studies on Alcohol and Drugs*. 2008; 69(4):481–90. [PubMed: 18612562]
- Wechsler H, Nelson TE, Lee JE, Seibring M, Lewis C, Keeling RP. Perception and reality: A national evaluation of social norms marketing interventions to reduce college students' heavy alcohol use. *Journal of Studies on Alcohol and Drugs*. 2003; 64(4):484–94.
- Werch CE, Pappas DM, Carlson JM, DiClemente CC, Chally PS, Sinder JA. Results of a social norm intervention to prevent binge drinking among first-year residential college students. *Journal of American College Health*. 2000; 49(2):85–92. [PubMed: 11016132]




Figure 1.
Example misperception function.




Figure 2. HED fraction as a function of simple one-variable treatments. Upper panel shows the effect of reducing misperception from 0.25 to 0.05. Lower panel shows the effect of lowering wetness from 0.30 to 0.00.




Figure 3. HED fractions as a function of changing underage wetness and misperception. The upper panel shows a campus with moderate wetness ($w=0.30$) among the legal age students; the lower panel shows a wet campus.

Table 1

Bounds for the rate parameters as estimated from SNMRP data.

Parameter Name	Value at w=0	Value at w=1
s12	0.0170	19.1313
s42	4.2113	3.9486
r31	4.6484	4.6133
r23	0.2538	0.3971
r24	1.2860	6.8722
rR42	6.3111	6.2014
r43	1.5545	1.5776
r21	0.58-0	0.1807
r12	0.5006	8.0899
r24	2.6021	4.3106